Winter 2019 Math 106
Topics in Applied Mathematics
Data-driven Uncertainty Quantification

Yoonsang Lee (yoonsang.lee@dartmouth.edu)

Lecture 10: Hilbert Space

10.1 Function Space

- A function space is a set of functions $\mathcal{F}$ that has some structure.
- Often a nonparametric density estimation or a function approximation is chosen to lie in some function space, where the assumed structure is exploited by algorithms and theoretical analysis.

Let V be a vector space over $\mathbb{R}$. A **norm** is a mapping $\|\cdot\| : V \to [0, \infty)$ that satisfies

1. $\|x + y\| \leq \|x\| + \|y\|$.
2. $\|ax\| = |a|\|x\|$ for all a $\in \mathbb{R}$.
3. $\|x\| = 0$ implies $x = 0$.

A vector space equipped with a norm is called a **normed vector space**.

Example. $V = \mathbb{R}^k$ with $\|x\| = \sqrt{\sum_i x_i^2}$.

10.1 Function Space

- A sequence $x_1, x_2, \ldots$ is said to converge to x if $\|x_n - x\| \to 0$ as $n \to \infty$.
- A sequence $x_1, x_2, \ldots$ in a normed space is a **Cauchy sequence** if $\|x_m - x_n\| \to 0$ as $m, n \to \infty$.
- The space is **complete** if every Cauchy sequence converges to a limit.
- A complete, normed vector space is called a **Banach space**.

Example. $L^p([0,1])$ spaces, $1 \leq p \leq \infty$

$$\{f(x) : \int |f^p(x)|^p dx < \infty\}.$$

10.2 Hilbert Space

- An **inner product** is a mapping $< \cdot, \cdot >: V \times V \to \mathbb{R}$ that satisfies, for all $x, y, z \in V$ and $a \in \mathbb{R}$:
  1. $\langle x, y \rangle \geq 0$ and $\langle x, x >= 0$ if and only if $x = 0$
  2. $\langle x, y + z >= \langle x, x > + \langle x, z >$
  3. $\langle x, ay >= a\langle x, y \rangle$
  4. $\langle x, y \rangle =< y, x >$

  Example. $V = \mathbb{R}^k$ with $\langle x, y \rangle = \sum_i x_i y_i$.
  Example. $V = L^2([0, 1])$ with $\langle f, g \rangle = \int f(x)g(x)dx$.
- x and y are **orthogonal** if $\langle x, y \rangle = 0$
- **Cauchy-Schwartz inequality**

$$|\langle x, y \rangle| \leq \|x\| \|y\|.$$

Example. $\int f(x)g(x)dx \leq (\int f^2 dx)^{1/2} (\int g^2 dx)^{1/2}$

10.2 Hilbert Space

- An inner product space is a normed space with the norm $\|x\| = \langle x, x \rangle$.
- Parallelogram property

$$\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2).$$

- A **Hilbert space**, H, is a complete, inner product vector space.
- Given a set $S \subset H$ of a normed linear space and some point b outside of S, the distance between b and S is defined as

$$d(b, S) = \inf_{x \in S} \|x - b\|.$$

- Note that in general there is no guarantee that there exists a point $u \in S$ such that $d(b, S) = \|u - b\|$ (this is why we have inf instead of min).

10.2 Hilbert Space

Theorem. A set $S \subset H$ is called **closed** if every convergent sequence $\{x_n\}$ in S converges to an element of S. If S is a closed linear space of a Hilbert space H and b is an element of H, then there exists $u \in S$ such that $\|u - b\| = d(b, S)$.

Idea of Proof.

- There exists a sequence $\{u_n\} \in S$ such that $\|u_n - b\| \to d(b, S)$ as $n \to \infty$.
- This does not mean that $\{u_n\}$ has a limit **in** S in general.
- From

$$\|\frac{1}{2}(b-u_m)\|^2+\|\frac{1}{2}(b-u_n)\|^2 = \frac{1}{2}\|b-\frac{1}{2}(u_n+u_m)\|^2+\frac{1}{8}\|u_n-u_m\|^2,$$

  $\|u_n - u_m\| \to 0$ and thus $\{u_n\}$ is a Cauchy sequence.
- From the definition of the Hilbert space, there exists $u \in S$ such that $d(b, S) = \|u - b\|$.

10.2 Hilbert Space

- For a closed subspace S of H and $x \in S$, $\hat{x}$ such that $d(x, S) = \|\hat{x} - x\|$ is called the **closest point** of x in S or the **projection** of x onto S.

Theorem. Let S be a closed linear subspace of H, let x be any element of S, b any element of V, and $\hat{b}$ the project of b onto S. Then

$$\langle x - \hat{b}, b - \hat{b} \rangle = 0.$$

Proof. If $x = \hat{b}$, we are done. Otherwise, set

$$\theta(x-\hat{b})-(b-\hat{b}) = \theta x+(1-\theta)\hat{b}-b = y-b \text{ where } y = \theta x+(1-\theta)\hat{b}.$$

Since y is in S and $\|y - b\| \geq \|\hat{b} - b\|$, we have

$$\|\theta(x-\hat{b})-(b-\hat{b})\|^2 = \theta^2\|x-\hat{b}\|^2-2\theta(x-\hat{b}, b-\hat{b})+\|b-\hat{b}\|^2 \geq \|b-\hat{b}\|^2.$$

Therefore, $\theta^2\|x - \hat{b}\|^2 - 2\theta\langle x - \hat{b}, b - \hat{b}\rangle \geq 0$ for all θ.

10.2 Hilbert Space

- For a closed subspace S of H and $x \in S$, $\hat{x}$ such that $d(x, S) = \|\hat{x} - x\|$ is called the **closest point** of x in S or the **projection** of x onto S.

Theorem. Let S be a closed linear subspace of H, let x be any element of S, b any element of V, and $\hat{b}$ the project of b onto S. Then

$$\langle x - \hat{b}, b - \hat{b} \rangle = 0.$$

Proof.
Therefore, $\theta^2 \|x - \hat{b}\|^2 - 2\theta \langle x - \hat{b}, b - \hat{b} \rangle \geq 0$ for all θ.
The left-hand side attains its minimum value when
$\theta = \langle x - \hat{b}, b - \hat{b} \rangle / \|x - \hat{b}\|^2$, in which case

$$-\langle x - \hat{b}, b - \hat{b} \rangle^2 / \|x - \hat{b}\|^2 \geq 0.$$

This implies

$$\langle x - \hat{b}, b - \hat{b} \rangle = 0.$$

10.2 Hilbert Space

- For a closed subspace S of H and $x \in S$, $\hat{x}$ such that $d(x, S) = \|\hat{x} - x\|$ is called the **closest point** of x in S or the **projection** of x onto S.

Theorem. Let S be a closed linear subspace of H, let x be any element of S, b any element of V, and $\hat{b}$ the project of b onto S. Then

$$\langle x - \hat{b}, b - \hat{b} \rangle = 0.$$

Proof.
Corollary. $b - \hat{b}$ is orthogonal to S.
Corollary. $\hat{b}$ is unique.

10.3 Parametric Regression

Example. For a given data set $\{(x_i, y_i), i = 1, 2, ..., m\}$, let we want a linear function fit to the data

$$y_i = ax_i + b.$$

In a matrix form, we are looking for a and b such that

$$\begin{pmatrix} x_1 & 1 \\ \vdots & \vdots \\ x_m & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix}$$

In a more general form, we are looking for a solution u to the following linear system

$$Au = v$$

where A is an $m \times n$ and $m > n$.

10.3 Parametric Regression

Example. (cont'd) If we assume that the column vectors of A are linearly independent, what can we say about the existence of the solution u?

- If $Au = v$ has a solution, then one can express v as a linear combination of $A_1, A_2, ..., A_n$ (the column vectors of A). That is, if v is not in the column space of A, there is no solution.
- The best we can represent about v is the projection of v into the column space of A, $\hat{v}$.
- Then, does $Au = \hat{v}$ has a solution? From the previous slides, we know

$$\langle A_1, \hat{v} - v \rangle = 0, \langle A_2, \hat{v} - v \rangle = 0, ..., \langle A_n, \hat{v} - v \rangle = 0.$$

That is, $A^T(Au - v) = A^T(\hat{v} - v) = 0$, i.e., $A^T Au = A^T v$. As $A^T A$ is invertible, we have

$$u = (A^T A)^{-1} A^T v, \text{the } \textbf{regression formula}!$$

10.4 Orthonormal Basis

Definition. An orthonormal basis of a Hilbert Space H is a family $\{e_k \in H\}_{k \in B}$ if it satisfies

1. $\langle e_k, e_j \rangle = 0$ for all $k \neq j$, $k, j \in B$.
2. $\|e_k\| = 1$ for all $k \in B$.
3. The linear span of $\{e_k\}$ is dense in H.

If the index set B is countable, the Hilbert space is called **separable**. That is, for any $u \in H$, u can be represented as

$$u = \sum_{i \in B} \beta_i e_i$$

for an orthonormal basis $\{e_i\}$.

Note. We will consider only separable Hilbert spaces.

- $\beta_i = \langle u, e_i \rangle$.
- $\|u\|^2 = \sum_{i \in B} \beta_i^2$.

10.4 Orthonormal Basis

Examples.

- $\{e^{2\pi inx}\}_{n=-\infty}^{\infty}$ is an orthogonal basis of $L^2[0,1]$.
- The Legendre polynomial is another orthogonal basis for $L^2[0,1]$.
- The Hermite polynomial is an orthogonal basis of $L^2(\mathbb{R})$.