

(1) (9 points) Let f be a function of three variables x , y and z . For each of the following assertions, select the correct ending.

(a) The integral $\int_0^{2t} \int_0^x \int_0^{xz} f(x, y, z) dy dz dx$ is...

- a constant.
- a function of t .
- a function of y .
- not well-defined.
- none of the above.

(b) The integral $\int_0^1 \int_x^0 \int_0^{xz} f(x, y, z) dy dx dz$ is...

- a constant.
- a function of x and z .
- a function of y .
- not well-defined.
- none of the above.

(c) The integral $\int_0^1 \int_0^2 f(x, y, z) dx dz$ is...

- a constant.
- a function of x and z .
- a function of y .
- not well-defined.
- none of the above.

(2) (7 points) Calculate the double integral

$$\iint_{\mathcal{R}} \left(\frac{x}{y} + \frac{y}{x} \right) dA$$

where $\mathcal{R} = [1, 4] \times [1, 2]$.

(3) (7 points) Evaluate the following integral:

$$\int_0^1 \int_{3y}^3 e^{x^2} dx dy.$$

(4) (7 points) Calculate the triple integral $\int_0^{\sqrt{\pi}} \int_0^x \int_0^{xz} x^2 \cos(y) dy dz dx$.

- (5) (8 points) Find the volume of the solid that lies below the surface $z^2 = 9x^2 + 9y^2$, above the xy -plane, and inside the cylinder $x^2 + y^2 = 2y$.

Hint: you may use without proof any of the following facts:

$$\int_0^{\frac{\pi}{2}} \sin^3 \theta \, d\theta = \frac{2}{3} \quad , \quad \int_0^{\pi} \sin^3 \theta \, d\theta = \frac{4}{3} \quad , \quad \int_0^{\frac{\pi}{2}} \sin^2 \theta \, d\theta = \frac{\pi}{4} \quad , \quad \int_0^{\pi} \sin^2 \theta \, d\theta = \frac{\pi}{2}.$$

(6) (7 points) The volume of the solid bounded by the surfaces with equations

$$y = x^2 + \frac{z^2}{4} \quad \text{and} \quad y = 5 - 4x^2 - z^2$$

is given by an integral of the form

$$\int_a^b \int_{u_1(z)}^{u_2(z)} \int_{x^2 + \frac{z^2}{4}}^{5 - 4x^2 - z^2} dy \, dx \, dz.$$

Determine $u_1(z)$, $u_2(z)$, a and b .

(7) (7 points) Let \mathcal{W} be the solid region lying:

- inside the cone $z = \sqrt{x^2 + y^2}$
- inside the sphere $x^2 + y^2 + z^2 = 5$
- outside the sphere $x^2 + y^2 + z^2 = 3$
- in the **second** octant: $x \leq 0, y \geq 0, z \geq 0$.

Convert the integral $\iiint_{\mathcal{W}} x \, dV$ into a triple integral in spherical coordinates.

Do not evaluate the integral.

(8) (8 points) Express the volume of the region bounded by the surfaces with equations

- $z = x^2 - 1$
- $x = y^2$
- $z = 0$

by a triple integral in the prescribed orders.

Do not evaluate the integrals.

(a) $dz dy dx$

(b) $dy dx dz$

- (9) (8 points) A lamina has the shape of a half-disk of radius 2, centered at the origin and lying in the upper-half plane. Its density function is $\delta(x, y) = k\sqrt{x^2 + y^2}$, where k is a constant.

The center of mass of the lamina has coordinates $(0, y_0)$. Determine y_0 .

This page is intentionally left blank. You can use it as scratch paper.

This page is intentionally left blank. You can use it as scratch paper.