

# Integral Equations

**Volterra** operator  $Ku(x) = \int_0^x k(x,y) u(y) dy$

'lower-triangular' kernel  


- $u - \lambda Ku = f$  has unique solution for any  $\lambda$ , continuous func.  $f$ .  
This is  $u = (1 + \lambda K + \lambda^2 K^2 + \dots) f$ . Neumann Series (converges)
- this tells you  $K$  has no eigenvalues
- The other way to solve a Volterra eqn. is to take derivs. via Leibniz's formula until you get an ODE, then solve that, with ICs that can be extracted from the  $x \rightarrow 0$  limit of the integral eqn.
- You can go backwards, i.e. given an ODE convert to Volterra eqn. Make sure you can do this for 1<sup>st</sup> order. (Don't do 2<sup>nd</sup> order since harder, see p.283)
- Volterra eqns arise in real-world situations where  $u(t)$  determined by history  $u(s)$  for  $s < t$ .

**Fredholm degenerate**

op.  $Ku(x) = \sum_{j=1}^N \alpha_j(x) \beta_j(y)$   $\{\alpha_j\}$  L.I. set  
 $\{\beta_j\}$  " "  
 $\Rightarrow K$  need not be symm.

- Eigenvalues are those of matrix  $A$  with entries  $A_{ij} = (\beta_i, \alpha_j)$ , plus an  $\infty$ -multiplicity zero eigenvalue.
- Eigenfuncs are  $\sum_{j=1}^N c_j \alpha_j(x)$  where  $\vec{c}$  is corresponding eigenvector of  $A$ , plus the set of all funcs orthogonal to all  $\{\beta_j\}$  forms the zero eigenspace
- $Ku - \lambda u = f$  has unique soln if  $\lambda \neq$  eigenvalue, which can be got from  $\sum_{j=1}^N \alpha_j(x) c_j - \lambda u(x) = f(x)$  (\*)  
But if  $\lambda = j^{\text{th}}$  eigenvalue then no soln. unless  $\vec{f} = \{f_i\}$   $f_i = (\beta_i, f)$  is in the range of  $A\vec{c} - \lambda\vec{c}$ , i.e.  $A\vec{c} - \lambda\vec{c} = \vec{f}$  consistent.
- $Ku = f$  has no soln. unless  $f$  is in  $\text{Span}\{\alpha_j\}$   $\leftarrow$  the range of  $K$ . when soln is nonunique in the zero eigenspace component.

**Fredholm symmetric**

op.  $Ku(x) = \int_1^b k(x,y) u(y) dy$

with  $k(y,x) = k(x,y)$  continuous.

so  $(Ku, v) = (u, Kv) \quad \forall u, v \in L^2$ .

- Eigenvalues  $\lambda_j$  real, tend to zero, a number of them.
- Eigenfunctions  $\phi_j$  orthogonal, complete in  $L^2$  ... means form a basis for  $L^2$ .

All your techniques from symmetric matrices work:

•  $Ku - \lambda u = f$       write  $u = \sum_{j=1}^{\infty} c_j \phi_j$  ,  $f = \sum_{j=1}^{\infty} f_j \phi_j$

gives  $c_j = \frac{f_j}{\lambda_j - \lambda}$       by orthogonality       $\uparrow$  you may need to compute.

$\hookrightarrow$  tells you: unique soln. if  $\lambda \neq$  eigenvalue

otherwise nonunique solution:  $u(x) = c \phi_j(x) + \sum_{i \neq j} \frac{f_i \phi_i(x)}{\lambda_i - \lambda}$

(if  $\lambda = \lambda_j$  for some  $j$ )       $\uparrow$  arbitrary.      if  $f_i = 0$ .

or no solution if  $f_j \neq 0$ .

• This all applies for  $\lambda = 0$  too.

•  $K$  operator can be written in spectral form  $K = \sum_{j=1}^{\infty} \lambda_j \underbrace{\phi_j(\phi_j, \cdot)}_{\text{projection onto } j^{\text{th}} \text{ eigenfunc.}}$

Equivalent to diagonalizing a symm. matrix.