


# Math 63, Winter 2010

## Homework set 6

### The Hilbert curve (1891).

The following picture is taken from the wikipedia entry for *Space-filling curve*.



This picture consists of six separate graphs, each depicting a curve in the plane. In fact, the six curves depicted here represent the first six functions in a sequence  $f_1, f_2, f_3, \dots$  with  $f_n: [0, 1] \rightarrow \mathbb{R}^2$ . Each picture shows, in red, a subset of the plane  $\mathbb{R}^2$  that is the *image*  $f_n([0, 1])$  for one of these functions. The particular sequence of functions depicted here was constructed by the mathematician David Hilbert in 1891, inspired by a similar idea of Guiseppe Peano from the year before.

Let's first look at the picture in the upper left hand corner. It depicts the image of a function  $f_1: [0, 1] \rightarrow \mathbb{R}^2$ . To describe the nature of the function  $f_1$ , we divide the interval  $[0, 1]$  into four equal (but not disjoint) parts

$$[0, 1] = [0, \frac{1}{4}] \cup [\frac{1}{4}, \frac{2}{4}] \cup [\frac{2}{4}, \frac{3}{4}] \cup [\frac{3}{4}, 1] = I_1 \cup I_2 \cup I_3 \cup I_4.$$

We also divide the square  $[0, 1] \times [0, 1] \subset \mathbb{R}^2$  into 4 equal parts:

$$\begin{aligned} [0, 1] \times [0, 1] &= [0, \frac{1}{2}] \times [0, \frac{1}{2}] \cup [0, \frac{1}{2}] \times [\frac{1}{2}, 1] \cup [\frac{1}{2}, 1] \times [\frac{1}{2}, 1] \cup [\frac{1}{2}, 1] \times [0, \frac{1}{2}] \\ &= S_1 \cup S_2 \cup S_3 \cup S_4. \end{aligned}$$

Then the crucial property of the function  $f_1$  is that it maps each interval  $I_p$  to the square  $S_p$ ,

$$f_1(I_p) \subset S_p \text{ for } p = 1, 2, 3, 4.$$

(Make sure you see how these facts relate to the image in the first picture.)

Moreover, if you look at all the five pictures of  $f_2, f_3, f_4, f_5, f_6$  that come after  $f_1$ , you may see that they have the same property:

$$f_n(I_p) \subset S_p, \text{ for } p = 1, 2, 3, 4, \text{ and for every } n \geq 1.$$

The curves  $f_n$  become more and more convoluted, but remain subordinate to this rule imposed by the division of the interval  $[0, 1]$  and the square  $[0, 1] \times [0, 1]$  into four equal parts.

In general, for every value of  $n = 1, 2, 3, \dots$ , there exists a subdivision of the interval  $[0, 1]$  into  $4^n$  closed intervals of size  $1/4^n$ ,

$$[0, 1] = \bigcup_{p=1}^{4^n} I_p^n.$$

The explicit formula for the interval  $I_p^n$  is

$$I_p^n = \left[ \frac{p-1}{4^n}, \frac{p}{4^n} \right].$$

Likewise, the square  $[0, 1] \times [0, 1] \subset \mathbb{R}^2$  can be sub-divided into  $4^n$  smaller squares with sides of length  $1/2^n$ . Let us assume that we label this set of squares somehow as  $S_1^n, S_2^n, \dots, S_{4^n}^n$ , so that

$$[0, 1] \times [0, 1] = \bigcup_{p=1}^{4^n} S_p^n,$$

where each  $S_p^n$  is of the form

$$S_p^n = \left[ \frac{q-1}{2^n}, \frac{q}{2^n} \right] \times \left[ \frac{r-1}{2^n}, \frac{r}{2^n} \right],$$

for some choice of integers  $q, r$  between 1 and  $2^n$ .

Hilbert proved (and you may take this for granted) that it is possible to label the squares  $S_p^n$  in such a way that there exists a *continuous* function  $f_n: [0, 1] \rightarrow \mathbb{R}^2$  which satisfies

$$f_n(I_p^n) \subset S_p^n \text{ for } p = 1, 2, \dots, 4^n.$$

Moreover, all of the later functions  $f_m$  in the sequence (i.e.,  $m > n$ ) also respect this particular subdivision into  $4^n$  parts,

$$f_m(I_p^n) \subset S_p^n, \text{ for all } m \geq n, p = 1, 2, \dots, 4^n.$$

Before you try to solve the following problems, identify the squares  $S_p^n$  in the six pictures and try to understand how the properties of  $f_n$  mentioned here are reflected in the graphs.

1. Prove that the sequence of functions  $f_1, f_2, f_3, \dots$  has a limit  $f = \lim_{n \rightarrow \infty} f_n$ , and that the limit is a *continuous* function  $f: [0, 1] \rightarrow \mathbb{R}^2$ . (Hint: Prove that it is a Cauchy sequence in the metric space  $\mathcal{F}$  of continuous functions from the closed interval  $E = [0, 1]$  to Euclidean space  $E' = \mathbb{R}^2$ .)
2. Prove that every point  $(x, y) \in [0, 1] \times [0, 1]$  occurs in the image  $f([0, 1])$  of the limit function. (The limit  $f$  is called the *Hilbert curve* and it is an example of a *space-filling curve*.)