

LECTURE OUTLINE
Integration by Parts

Professor Leibon

Math 8

Sep 24, 2004

Goal

Chain Rule, eluR niahC
Product Rule, eluR tcudorP
Trig Inverses

Reversing the Chain Rule

The chain rule assures us that

$$\frac{d(f(u))}{dx} = \frac{df}{du}(u) \frac{du}{dx},$$

hence we find the elur niahc

$$\int \frac{df}{du}(u) \frac{du}{dx} dx = f(u(x)) + C.$$

u-substitution

For pronunciation purposes, we express the elur niahc as

$$\int h(u) \frac{du}{dx} dx = \int h(u) du$$

where $f(u) = \int h(u) du$, and call its **use** *u*-substitution.

Example 1

Use u -substitution to find

$$\int \frac{x}{1+x^2} dx.$$

Reversing the Product Rule

The product rule assures us that

$$(uv)' = u'v + uv',$$

hence we find

$$\int uv' dx + \int u'v dx = uv + C.$$

Integration by Parts

For pronunciation purposes, we write the formula as

$$\int uv' dx = uv - \int u'v dx$$

and call its use *integration by parts*.


Example 2

Use integration by parts to find

$$\int \ln(x) dx.$$

$$\tan^{-1}$$

Let $\tan^{-1}(x)$ be a continuous inverse of $\tan(x)$,
when $\tan(x)$ has been restricted to the interval
 $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.



Derivative of \tan^{-1}

Prove that

$$\frac{d}{dx} \tan^{-1}(x) = \frac{1}{1+x^2},$$

and hence that

$$\int \frac{1}{1+x^2} dx = \tan^{-1}(x) + C.$$

Example 3

Find

$$\int \tan^{-1}(x) dx$$

Similarly for

$$\begin{aligned}\frac{d}{dx} \sin^{-1}(x) &= \frac{1}{\sqrt{1-x^2}} & \frac{d}{dx} \cos^{-1}(x) &= \frac{-1}{\sqrt{1-x^2}} \\ \frac{d}{dx} \tan^{-1}(x) &= \frac{1}{1+x^2} & \frac{d}{dx} \cot^{-1}(x) &= \frac{-1}{1+x^2} \\ \frac{d}{dx} \sec^{-1}(x) &= \frac{1}{x\sqrt{1-x^2}} & \frac{d}{dx} \csc^{-1}(x) &= \frac{-1}{x\sqrt{1-x^2}}\end{aligned}$$

Example 4

Find the area of the region
bounded by $y = \frac{\pi^2}{4}e^{(-x+\pi)} \sin(x)$
and $y = x^2e^x$ and the lines $x = 0$,
 $x = \frac{\pi}{2}$.